

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : C07H 21/02, 21/04, C07K 9/00 A61K 37/00, C12N 1/21, 5/10 C12N 15/70, C12P 21/00		A1	(11) International Publication Number: <b>WO 94/10187</b> (43) International Publication Date: 11 May 1994 (11.05.94)
(21) International Application Number: PCT/US93/10307 (22) International Filing Date: 26 October 1993 (26.10.93) (30) Priority data: 07/968,953 30 October 1992 (30.10.92) US 07/968,954 30 October 1992 (30.10.92) US (60) Parent Applications or Grants (63) Related by Continuation US 07/968,953 (CIP) Filed on 30 October 1992 (30.10.92) US 07/968,954 (CIP) Filed on 30 October 1992 (30.10.92) (71) Applicants (for all designated States except US): HSC RESEARCH AND DEVELOPMENT LIMITED PARTNERSHIP [CA/CA]; 88 Elm Street, Toronto, Ontario M5G 1X8 (CA). MEMORIAL SLOAN-KETTERING CANCER CENTER [US/US]; 1275 York Avenue, New York, NY 10021 (US). CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS [ES/ES]; Velazquez, 144, E-28006 Madrid (ES).		(72) Inventors; and (75) Inventors/Applicants (for US only): LETARTE, Michelle [CA/CA]; 35 Howland Road, Toronto, Ontario M4K 2Z7 (CA). MASSAGUE, Joan [US/US]; 1233 York Avenue, New York, NY 10021 (US). BERNABEU, Carmelo [ES/ES]; Centro Investigaciones Biologicas, Velazquez, 144, E-28006 Madrid (ES). CHEIFETZ, Sela [CA/CA]; 88 New Seabury Drive, Concord, Ontario L4K 2R8 (CA). (74) Agents: KONSKI, Antoinette, F. et al.; Campbell & Flores, 4370 La Jolla Village Drive, Suite 700, San Diego, CA 92122 (US). (81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
Published with international search report <b>AMIL</b> INFORMATION SERVICES P O BOX 405, CORTE MADERA, CA 94976-0405 (415) 927-0340 • FAX (415) 927-7250			
(54) Title: COMPOSITIONS AND METHODS FOR MODIFYING THE REGULATORY ACTIVITY OF TGF- $\beta$			
(57) Abstract <p>This invention provides a novel purified TGF-<math>\beta</math> binding glycoprotein, endoglin, and isolated nucleic acid molecules that encode amino acid sequences corresponding to the TGF-<math>\beta</math>-binding glycoprotein. Also provided is soluble endoglin-derived polypeptide, and fragments thereof. A pharmaceutical composition which comprises the purified endoglin-derived polypeptide or produced recombinantly methods and a pharmaceutically acceptable carrier is further provided as well as methods of treating patients by administering to the patient the pharmaceutical compositions of this invention.</p>			

BEST AVAILABLE COPY

62A1  
ATTORNEY DOCKET NUMBER: 10177-216-999  
SERIAL NUMBER: 09/590,002  
REFERENCE: B107

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : C07H 21/02, 21/04, C07K 9/00 A61K 37/00, C12N 1/21, 5/10 C12N 15/70, C12P 21/00		A1	(11) International Publication Number: WO 94/10187 (43) International Publication Date: 11 May 1994 (11.05.94)
(21) International Application Number: PCT/US93/10307 (22) International Filing Date: 26 October 1993 (26.10.93) (30) Priority data: 07/968,953 30 October 1992 (30.10.92) US 07/968,954 30 October 1992 (30.10.92) US (60) Parent Applications or Grants (63) Related by Continuation US 07/968,953 (CIP) Filed on 30 October 1992 (30.10.92) US 07/968,954 (CIP) Filed on 30 October 1992 (30.10.92) (71) Applicants (for all designated States except US): HSC RESEARCH AND DEVELOPMENT LIMITED PARTNERSHIP [CA/CA]; 88 Elm Street, Toronto, Ontario M5G 1X8 (CA). MEMORIAL SLOAN-KETTERING CANCER CENTER [US/US]; 1275 York Avenue, New York, NY 10021 (US). CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS [ES/ES]; Velazquez, 144, E-28006 Madrid (ES).		(72) Inventors; and (75) Inventors/Applicants (for US only) : LETARTE, Michelle [CA/CA]; 35 Howland Road, Toronto, Ontario M4K 2Z7 (CA). MASSAGUE, Joan [US/US]; 1233 York Avenue, New York, NY 10021 (US). BERNABEU, Carmelo [ES/ES]; Centro Investigaciones Biologicas, Velazquez, 144, E-28006 Madrid (ES). CHEIFETZ, Sela [CA/CA]; 88 New Seabury Drive, Concord, Ontario L4K 2R8 (CA). (74) Agents: KONSKI, Antoinette, F. et al.; Campbell & Flores, 4370 La Jolla Village Drive, Suite 700, San Diego, CA 92122 (US). (81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
		Publ. Int. Information search report <b>AMI</b> INFORMATION SERVICES P O BOX 405, CORTE MADERA, CA 94976-0405 (415) 927-0340 • FAX (415) 927-7250	
(54) Title: COMPOSITIONS AND METHODS FOR MODIFYING THE REGULATORY ACTIVITY OF TGF- $\beta$			
(57) Abstract <p>This invention provides a novel purified TGF-<math>\beta</math> binding glycoprotein, endoglin, and isolated nucleic acid molecules that encode amino acid sequences corresponding to the TGF-<math>\beta</math>-binding glycoprotein. Also provided is soluble endoglin-derived polypeptide, and fragments thereof. A pharmaceutical composition which comprises the purified endoglin-derived polypeptide or produced recombinantly methods and a pharmaceutically acceptable carrier is further provided as well as methods of treating patients by administering to the patient the pharmaceutical compositions of this invention.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

COMPOSITIONS AND METHODS FOR MODIFYING THE REGULATORY  
ACTIVITY OF TGF- $\beta$

FIELD OF THE INVENTION

The present invention relates to cell biology and  
5 to methods of modifying the biological activity of cell  
regulatory factors. More specifically, the present  
invention relates to a novel TGF- $\beta$ -binding glycoprotein.

Throughout this application various publications  
are referenced within parentheses. The disclosures of  
10 these publications in their entireties are hereby  
incorporated by reference to more fully describe the state  
of the art to which this invention pertains.

BACKGROUND OF THE INVENTION

Glycoproteins, in which one or more carbohydrate  
15 units have been attached covalently to the protein by  
posttranslational processing are widely distributed.  
Several secretory proteins, including the immunoglobulins,  
are glycoproteins, as are most components of plasma  
membranes such as cell membrane receptors, where the  
20 carbohydrates can be involved in cell-to-cell adhesion.

Transforming growth factor  $\beta$  (TGF- $\beta$ ) refers to a  
family of multi-functional cell regulatory factors produced  
in various forms by many cell types (for review see Sporn  
et al., J. Cell Biol. 105:1039 (1987)). Five distinct  
25 isoforms of TGF- $\beta$  have been identified. TGF- $\beta$ 1 and TGF- $\beta$ 2  
have been characterized in detail. TGF- $\beta$  is the subject of  
U.S. Patent Nos. 4,863,899; 4,816,561 and 4,742,003 which  
are incorporated herein by reference. TGF- $\beta$  binds to cell  
surface receptors present on various types of cells and is  
30 known to potentiate or inhibit the response of most cells  
to other growth factors, depending on the cell type. TGF- $\beta$   
also regulates differentiation of some cell types, either  
promoting or inhibiting proliferation of the cell. Another

marked effect of TGF- $\beta$  is the promotion of cellular production of extracellular matrix proteins and their receptors (for a review see Keski-Oja et al., J. Cell Biochem. 33:95 (1987); Massague, Cell 49:437 (1987);

5 Roberts and Sporn, "Peptides Growth Factors and Their Receptors", Springer-Verlag (1989)).

Notwithstanding the beneficial and essential cell regulatory functions served, TGF- $\beta$  regulatory activity can prove detrimental to its host organism. For example,

10 whereas growth and proliferation of mesenchymal cells is stimulated by TGF- $\beta$ , some tumor cells may also be stimulated, using TGF- $\beta$  as an autocrine growth factor. In other cases the inhibition of cell proliferation by TGF- $\beta$  similarly proves detrimental to its host organism. An

15 example would be the prevention of new cell growth to assist in repair of tissue damage. The stimulation of extracellular matrix production by TGF- $\beta$  is essential for wound healing. However, in some cases, the TGF- $\beta$  response is uncontrolled and an excessive accumulation of

20 extracellular matrix results. An example of excessive accumulation of extracellular matrix is the "internal" scarring that occurs in the pathology glomerulonephritis and dermal scar tissue formation.

25 The transforming growth factor- $\beta$  receptor system in most mesenchymal and epithelial cells consists of several components (Massague, J. Ann. Rev. Cell Biol. 6:597 (1990); Lin, H.Y. et al., Cell 68:775 (1992); Georgi, L.L. et al., Cell 61:635 (1990); Mathews, L.S. et al., Cell

30 65:973 (1991); Attisano, L. et al., Cell 68:97 (1992); Lopez-Casillas et al., Cell 67:785 (1991) and Wang et al., Cell 67:796 (1991)), one of which is betaglycan, a membrane-anchored proteoglycan. In addition to betaglycan, the TGF- $\beta$  receptor system in most mesenchymal and

35 epithelial cells consists of the type I receptor, a 53-kDa glycoprotein whose structure has not been determined yet,

and the type II receptor, which belongs to the protein serine/threonine kinase receptor family. Additional cell surface TGF- $\beta$ -binding proteins, some of which have a more restricted distribution, have also been described.

- 5                   Thus, a need exists to develop compounds that can modify the effects of cell regulatory factors such as TGF- $\beta$ . The present invention satisfies this need and provides related advantages as well.

#### SUMMARY OF THE INVENTION

- 10                   The present invention provides a novel purified TGF- $\beta$ -binding glycoprotein. The protein, endoglin, is expressed at high levels on human vascular endothelial cells.

- Further provided by the present invention are  
15 methods of treating pathologic conditions mediated by TGF- $\beta$  regulatory activity by contacting the TGF- $\beta$  with an effective amount of purified endoglin-derived polypeptide or any fragment thereof having the ability to bind TGF- $\beta$ . Thus, intact, native endoglin and soluble fragments thereof  
20 are useful in these methods. This invention provides a method of preparing and purifying full length and soluble endoglin-derived polypeptide. Isolated nucleic acids encoding the novel TGF- $\beta$ -binding glycoprotein and soluble endoglin-derived polypeptides are also provided, as well as  
25 vectors containing the nucleic acids and recombinant host cells transformed with such vectors.

#### BRIEF DESCRIPTION OF THE FIGURES

- Figure 1 shows the domain structures of betaglycan and endoglin. Shown is a schematic  
30 representation highlighting regions of similarity between the linear sequences of betaglycan, an 853-amino acid

transmembrane proteoglycan, and endoglin, a disulfide-linked transmembrane protein composed of two identical subunits of 633 amino acids each. The transmembrane and short cytoplasmic regions (dark shaded box) of endoglin have a high level of sequence similarity to the corresponding regions of betaglycan. Two regions of weaker similarity are detected in the ectodomains of these proteins (light shaded boxes). Numbers represent the percent amino acid sequence similarity between the indicated domains of betaglycan and endoglin. Closed ovals represent positions of cysteine residues. Two putative sites for glycosaminoglycan chain attachment in betaglycan are indicated.

Figure 2 shows cell surface TGF- $\beta$ 1-binding proteins expressed by HUVEC. Near confluent cultures of HUVECs were affinity-labeled by incubation with 100 pM  $^{125}$ I-TGF- $\beta$ 1 followed by chemical cross-linking with 0.16 mM disuccinimidyl suberate. A) Triton X-100 extracts of affinity-labeled HUVEC were resolved on SDS-PAGE gels under reducing (R) or nonreducing (NR) conditions. Lane C contains extract from cells affinity-labeled in the presence of excess unlabeled TGF- $\beta$ 1. The migration position of TGF- $\beta$  receptors I (RI) and II (RII) are indicated. Arrow, the major affinity-labeled proteins of 180 kDa and higher molecular mass apparent on nonreducing gels. Arrowhead, the affinity-labeled proteins of 110-120 kDa seen on reducing gels. B) Detergent extracts of affinity-labeled HUVEC were resolved under nonreducing conditions on a first gel that was then resolved under reducing conditions in the second dimension as described in Cheifetz and Massague, J. Biol. Chem. 266:20767-20772 (1991), incorporated herein by reference. The 110-120-kDa labeled species migrating off-the-diagonal are indicated (arrowheads).

Figure 3 shows specific immunoprecipitation of TGF- $\beta$ 1-endoglin complexes. HUVECs were affinity-labeled with 100 pM  $^{125}$ I-TGF- $\beta$ 1 as described in Figure 2. A) Detergent extracts of affinity-labeled cells were incubated with mAb 44G4 and immune complexes were collected on protein G-Sepharose. After washes, equal aliquots of the samples were analyzed under reducing (R) or non-reducing (NR) conditions by SDS-PAGE (5-8% polyacrylamide gradient gels). B) Affinity-labeled HUVEC lysates were maximally depleted of endoglin by two successive 45 min incubations at 4°C with 100  $\mu$ l of 44G4-IgG-Sepharose. S) supernatant after second immunoprecipitation. I) the first 44G4 immunoprecipitation which contained 83% of the endoglin. T) corresponding amount of total extract used for the depletion experiment. All samples were analyzed under nonreducing conditions on SDS-PAGE with the exception of I<sub>R</sub>, which was run under reducing conditions. The migration positions of TGF- $\beta$  receptor II (RII), and endoglin monomer, dimer, and oligomer are indicated.

Figure 4 shows that endoglin transiently expressed in COS-M6 cells binds TGF- $\beta$ 1. COS-M6 cells were transfected with a cDNA encoding full-length L-endoglin (Endoglin) or control vector (C). Cells were affinity-labeled with 150 pM  $^{125}$ I-TGF- $\beta$ 1 and the detergent extracts incubated with mAb 44G4 followed by protein G-Sepharose. Immunoprecipitated proteins were analyzed by SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by autoradiography.

Figure 5 shows the specificity of endoglin for TGF- $\beta$  isoforms assessed in COS cell transfectants and in HUVEC. A) COS-M6 cells transfected with endoglin vector were affinity-labeled with 150 pM  $^{125}$ I-TGF- $\beta$ 1 alone or in the presence of 1 or 10 nM unlabeled TGF- $\beta$ 1, - $\beta$ 2 or - $\beta$ 3. B) HUVEC were affinity-labeled with 100 pM  $^{125}$ I-TGF- $\beta$ 1 alone or in the presence of 5 nM unlabeled TGF- $\beta$ 1 or TGF- $\beta$ 2.



Lysates from these cells were immunoprecipitated with MAb 44G4. Immunoprecipitates were fractionated under reducing conditions on SDS-PAGE gels. The region of the gels containing monomeric endoglin is shown along with the migration position of 100-kDa marker.

Figure 6 shows the restriction enzyme map of vector pcNeoSolEND with the 1.7 kb endoglin cDNA insert.

Figure 7 shows the partial nucleotide and predicted amino acid sequences of isolated S-endoglin cDNA. Nucleotides are numbered on the right. Amino acids are numbered on the left. The predicted signal sequence (nucleotides 283-357) and transmembrane region (nucleotides 2042-2116) are boldfaced. The 135 bp insert (underlined) contains splicing consensus sequences of donor/acceptor sites (GT, AG at positions 2134 and 2267) and branch point of lariat (CTGAC at position 2234). Nucleotides 372-2021 and 2322-3073 were found to be identical to the corresponding cDNA sequence of endothelial endoglin (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990) and Table 1).

Figure 8 shows the analysis of the cytoplasmic region which reveals the existence of two different forms of endoglin. A. Diagram of the cDNA coding for the cytoplasmic regions of the two alternative forms of endoglin. Only the region corresponding to the 3' end containing the cytoplasmic domain is depicted. The isolated endoglin cDNA (S-endoglin) contains a 135 bp insert not present in the previously described sequence (L-endoglin) (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)). The sequence of the additional 135 bp insert is shown in Figure 7. The position of the stop codons and the corresponding translated protein sequences (thick bar), are indicated. B. Alignment of the cytoplasmic and transmembrane domains of S-endoglin with

the corresponding sequences of L-endoglin and human betaglycan, Morén, A. et al., Biochem. Biophys. Res. Commun. 189:356 (1992). Boxes contain identical sequences. Numbers indicate the position of the first amino acid in the whole sequence. Two major regions of identity were found. The first region (73% identity) involves residues 587-617 of S- and L-endoglin and residues 780-810 of betaglycan. The second region (74% identity), involves residues 634-660 of L-endoglin and residues 823-849 of betaglycan. The transmembrane region of S-endoglin is boldfaced. Dashes have been inserted for purposes of alignment. Asterisks indicate the last residue of the protein.

Figure 9 shows the expression of L-endoglin and S-endoglin in transfectant cells. Mouse fibroblasts were transfected with either L-endoglin or S-endoglin cDNA and the expression of the endoglin molecule analyzed. A. Analysis by cytofluorometry of the endoglin present at the cell surface. After trypsinization, cells were stained for indirect immunofluorescence with the monoclonal antibody 8E11 (anti-endoglin). A control staining of endoglin mock transfectants is also shown. B. Immunoprecipitation analysis. Cells were metabolically labeled with [<sup>35</sup>S]methionine, lysed and immunoprecipitated with 44G4 (anti-endoglin) or HC1/1 (anti CD11c) monoclonal antibody. Monoclonal antibody HC1/1 was included as a negative control. Samples were electrophoresed on a 6-12% acrylamide gradient gel under nonreducing conditions. C. Immunoblotting analysis. Mock (L cells), L-endoglin (L-Endo) and S-endoglin (S-Endo) cDNA transfected mouse fibroblasts and PMA-treated U937 cells were lysed in Triton X-100, and the insoluble material removed by centrifugation. Proteins contained in the supernatant were electrophoresed on a 6% acrylamide gel under non-reducing conditions and transferred to nitrocellulose membranes. Immunodetection of endoglin was carried out with 44G4

(anti-endoglin) monoclonal antibody using a chemiluminiscent assay. HCl/1 (anti-CD11c) and X63 monoclonal antibody were used as negative controls. D. Immunoprecipitation analysis of cell surface labeled  
5 endoglin. Mock, L-endoglin (L-Endo) and S-endoglin (S-Endo) transfected mouse fibroblasts were <sup>125</sup>I-labeled, lysed, and immunoprecipitated with 44G4 monoclonal antibody (anti-endoglin). Samples were electrophoresed on a 10% acrylamide gel under either reducing (R) or nonreducing  
10 conditions (NR).

Figure 10 shows the detection of L-endoglin and S-endoglin transcripts by PCR amplification. Samples of total RNA from placenta, PMA treated HL-60 cells or PMA-treated U937 cells were incubated either in the presence  
15 (+RT) or in the absence (-RT) of reverse transcriptase. The generated cDNA samples, together with cDNA from S-endoglin (S-Endo) or L-endoglin (L-Endo) clones in pUC13 and cDNA from an endothelial cell library, were used for PCR amplification in the presence of oligonucleotides #14  
20 and #15 specific for S-endoglin (panel A), or oligonucleotides #12 and #11 common to both L-endoglin and S-endoglin (panel B). No amplification was observed when the RT reaction was omitted, excluding the possibility of DNA contaminating the RNA samples. Additional bands below  
25 the specific amplified fragments of L and S endoglin probably represent primer-dimer artifacts.

Figure 11 shows that both S-endoglin and L-endoglin bind TGF-β1. Confluent cultures of S-endoglin (L+-S) and L-endoglin (L+-L) transfectants were affinity  
30 labeled by incubation with 100 pM <sup>125</sup>I-TGF-β1 alone or in the presence of 4 nM unlabeled TGF-β1, followed by chemical crosslinking with disuccinimidyl suberate. All samples bound specifically to TGF-β1, as revealed by a fourfold ratio between the cpm bound in the absence versus presence  
35 of cold competing ligand; the parental L cells and mock

transfectants bound on average 60,000 specific cpm, while the endoglin transfectants bound on average 110,000 specific cpm. Immunoprecipitates of the L+-S transfectant with 44G4-IgG Sepharose contained 7700 cpm on average  
5 versus 700 cpm with control IgG-Sepharose; immunoprecipitates of the L+-L transfectant contained 4300 cpm versus 630 cpm for control. These immunoprecipitates were run on a 6-9% acrylamide gradient gel in reducing (R) or nonreducing (NR) conditions. The positions of the endoglin  
10 monomer, dimer and oligomer and of molecular weight markers are indicated.

Figure 12 shows the multiple cloning sites of pcDNAI/Neo.

Figure 13 shows the amino acid and nucleotide  
15 sequence of "L-Endoglin."

Table 1

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
84

DETAILED DESCRIPTION OF THE INVENTION

Endoglin is a homodimeric membrane glycoprotein composed of disulfide-linked subunits. Human-derived endoglin has been shown to exist in at least two isoforms expressed from a human cDNA library, an "L-isoform," of about 90 kDa or an "S-isoform" of about 85 kDa, reduced. Human endoglin, purified from tissue is shown to be composed of two disulfide-linked subunits each of about 95 kDa. It is expressed in human pre-erythroblasts, macrophages, leukemic cells of the lymphoid and myeloid lineages and at higher levels in vascular endothelial cells. It is also abundant on the syncytiotrophoblast, the multinucleated placental layer which constitutes the interface with maternal blood and plays an important role in providing nutrient exchange and immunological protection of the fetus (Gougos et al., Inter. Immunol. 4:83-92 (1992)).

Endoglin was first identified on a pre-B leukemic cell line, by its reactivity with mAb 44G4 (Quackenbush and Letarte, J. Immunol. 134:1276-1285 (1985)). It is present at low levels on cells derived from childhood acute leukemia cases (ALL) with pre-B lymphoid and myeloid phenotype; it is absent from T-ALL (Gougos and Letarte, J. Immunol. 141:1925-1933 (1988b); Kreindler et al., Leukemia and Lymphoma 3:7-18 (1990)). Interestingly, the human endoglin gene is localized to chromosome 9q34-qter, likely within the region translocated to chromosome 22 in Philadelphia chromosome-positive leukemia (Fernandez-Ruiz et al., Cytogen. Cell Gen. 64:204-207 (1993)). In normal adult bone marrow, only 3-5% of mononuclear cells express endoglin and they bear a pro-erythroblast phenotype; pre-B and myeloid precursors do not show detectable levels of endoglin (Buhring et al., Leukemia 5:841-847 (1991)). Normal B and T lymphocytes and unstimulated monocytes do not express endoglin but an up-regulation is observed on

activated macrophages (Lastres et al., Eur. J. Immunol. 22:393-397 (1992)).

Endoglin expression is considerably increased in the endothelium of various pathological skin lesions where  
5 endothelial cell proliferation is known to occur (Westphal et al., J. Invest. Dermatol. 100:27-34 (1993)). In tumors, capillary endothelial cells undergoing active angiogenesis also show higher levels of endoglin than resting endothelium in adjacent tissue.

10 Purified human endoglin exists in various isoforms and is composed of two disulfide linked subunits of Mr=95,000; 90,000; or 85,000, and bears N- and O-linked oligosaccharides. The primary sequence of human endoglin is composed of a 25 amino acid signal sequence, an  
15 extracellular domain of about 561 amino acids, a single transmembrane region of 25 amino acids and a cytoplasmic tail of 14 to 47 amino acid residues (Gougos and Letarte, J. Biol. Chem. 265:8361-8364 (1990)).

A relationship between human endoglin and the  
20 TGF- $\beta$  receptor system was discovered with the molecular cloning of the rat TGF- $\beta$ -binding proteoglycan, betaglycan (also known as the type III TGF- $\beta$  receptor), which revealed that the transmembrane domain and the relatively short (43 amino acid) cytoplasmic tail of this protein were  
25 remarkably similar (71% amino acid sequence similarity and 63% amino acid identity) to the corresponding regions in endoglin (see Figure 1). The extracellular domains of these two proteins show limited homology in primary structure, and while endoglin is not a proteoglycan, it  
30 does contain N- and O-linked oligosaccharides.

Cloning of the TGF- $\beta$  receptor II (Lin et al., Cell 68:775-785 (1992)) revealed a functional transmembrane serine/threonine kinase which had previously been

identified as a polypeptide of 80 kd, when bound and chemically cross-linked to  $^{125}\text{I}$ -TGF- $\beta$  (Cheifetz et al., J. Biol. Chem. 265:20533-20538 (1990); Cheifetz and Massagué, J. Biol. Chem. 266:20767-20772 (1991); Massagué, J., Cell 5 69:1067-1070 (1992)). This receptor associates with the type I receptor (a 53 kd protein) to transduce the TGF- $\beta$  signals; it is proposed that receptor I needs receptor II to bind TGF- $\beta$  and that receptor II needs receptor I to mediate a signal (Laiho et al., J. Biol. Chem. 266:9108-10 9112 (1991); Laiho et al., J. Biol. Chem. 265:18518-18524 (1990); Wrana et al., Cell 71:1003-1014 (1992)). Transfection experiments with a recently cloned receptor I, also a serine/threonine kinase, further supports the view that receptor I must associate with receptor II to bind 15 TGF- $\beta$  (Ebner et al., Science 260:1344-1348 (1993)).

The present invention provides a soluble endoglin-derived human polypeptide that binds TGF- $\beta$ . The full-length soluble endoglin-derived polypeptide comprises a signal sequence that is cleaved during processing, the 20 561 amino acids of the extracellular domain of the mature endoglin polypeptide, an integral membrane protein, which consists of about 600 or 633 amino acids in total. Nucleic acid sequences encoding the human endoglin polypeptide are identified in Table 1 and Figure 13. The nucleic acid 25 sequences encoding the soluble endoglin-derived polypeptides are included within the sequences set forth in Table 1 (from about amino acid number 1 to about amino acid number 561) and Figure 13. Also provided by this invention is a vector having inserted therein the genomic DNA 30 molecule encoding endoglin. This vector was deposited under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure, with Coleccion Espanola de Cultivos Tipo (CECT), 46100 Burjasot 35 (Valencia), Spain on October 21, 1993, under CECT 4475. Accordingly, isolated genomic DNA encoding endoglin is



within the scope of this invention.

This invention also provides purified human endoglin polypeptides encoding two isoforms differing from each other in the cytoplasmic region. The "S-endoglin" has 14 amino acid residues in the cytoplasmic region and the "L-endoglin" has 47 amino acid residues in the cytoplasmic region. The 586 amino acids spanning the extracellular and transmembrane regions in the mature endoglin are identical.

As used herein, the term "purified" means that the molecule or compound is substantially free of contaminants normally associated with a native or natural environment. For example, the mature human proteins can be obtained from a number of methods. The methods available for the purification of membrane proteins include precipitation, gel filtration, ion-exchange, reverse-phase, and affinity chromatography. Other well-known methods are described in Deutscher et al., Guide to Protein Purification: Methods in Enzymology Vol. 182, (Academic Press 1990), which is incorporated herein by reference. Alternatively, a purified polypeptide of the present invention can also be obtained by well-known recombinant methods as described, for example, in Sambrook et al., Molecular Cloning: A Laboratory Manual 2d ed. (Cold Spring Harbor Laboratory 1989), also incorporated herein by reference. An example of this means for preparing soluble endoglin-derived polypeptide is to express nucleic acid encoding the soluble endoglin in a suitable host cell, such as a bacterial, yeast or mammalian cell, using methods well known in the art, and recovering the expressed soluble protein, again using methods well known in the art.

For the purpose of illustration, expression of soluble endoglin was achieved by excising a full length 2.3 kb endoglin cDNA fragment from pcEXV-L ENDO, Hind III linkers were added and, after digestion with Hind III, it

was subcloned into the multiple cloning site of pBluescript vector (Stratagene), followed by digestion with Tth 111I. (This enzyme cuts the coding region approximately 80 bp upstream of the transmembrane region.)

5 Synthetic, complementary oligomers were engineered to contain an in-frame stop codon and a Bam HI overhang, and were ligated to the linearized plasmid. After digestion with Hind III, and Bam HI, and purification of the 1.7 kb fragment, it was ligated to pCDNAI/Neo  
10 (Invitrogen, San Diego, Figure 12). After cloning, the clones were screened by restriction enzyme analysis for the correct orientation and were found to be in the correct orientation. Verification of the cDNA insert was done using T7 and SP6 sequencing primers to confirm the presence  
15 of the start and stop codons at the 5' and 3' ends of the insert.

5' End Primer

TAATACGACTCACTATAGGGAGACCCAAGCTTGGGGAATTCCGTGGACAGCATG

T7 PRIMER

HINDIII ECORI

Initiation

20 3' End Primer

AAGACCGTCTAGACGGATCCACTAG.....CTATAGTGTCACCTAAATG

BAMHI

SP6 PRIMER

To express the construct, CHO-kl cells (ATCC) were transfected with pCDNA/Neo to determine optimal  
25 conditions required to generate stable transfectants resistant to G418. Five (5) x 10<sup>6</sup> cells electroplated with 5 µg of DNA at a voltage of 300 volts and a time constant of 17.9 msec., using the 960 µF capacitator (Biorad Gene Pulse) yielded the most number of stable transfectants with  
30 minimal cell death.

The following flow chart illustrates a means to express nucleic acid encoding soluble endoglin.

16

## SOLUBLE ENDOGLIN

plasmid pcEXV-L ENDO  
Eco R1 digest

5 Isolate 2.3 kb fragment

Add Hind III linker & Hind III digest

10 Subclone into pBluescript  
MCS-Hind III  
& selection on X-gal

15 pBluescript-L Endo

20 Digest with Tth 111I to linearize

Add complementary oligomers creating  
stop codon & Bam HI overhang

25 TCTAGACG  
AGATCTGCCTAG

30 Hind III Digest

Isolate 1.7 kb Hind III-BamHI fragment

35 Ligate with dephosphorylated  
Hind III-Bam HI fragment of vector  
pcDNA1/Neo

40 Transform E. coli MC1061/P3  
selection on Tetracyclin & Ampicillin

The soluble polypeptide and biologically active fragments thereof can also be produced by chemical synthesis. Synthetic polypeptides can be produced using Applied Biosystems, Inc. Model 430A or 431A automatic polypeptide synthesizer and chemistry provided by the manufacturer. The soluble polypeptide can also be isolated directly from cells which have been transformed with the expression vectors described below in more detail.

As used herein, endoglin-derived polypeptide means a human polypeptide having the amino acid sequence substantially the same as the 633 amino acid sequence shown in Table 1, or the 658 amino acid sequence shown in Figure 13, or an active fragment thereof. As used herein the term "soluble endoglin-derived polypeptide" refers to a soluble, biologically active fragment of the human endoglin polypeptide expressed by the extracellular domain of the nucleic acid. As used herein, an "active fragment" or "biologically-active fragment" refers to any portion of an endoglin polypeptide that binds to TGF- $\beta$ . Methods of determining whether a polypeptide can bind TGF- $\beta$  are well known to those of skill in the art, for example, as set forth herein.

The invention also encompasses nucleic acid molecules which differ from that of the nucleic acid molecule shown in Table 1, e.g., the sequence shown in Figure 13, but which produce the same phenotypic effect. These altered, but phenotypically equivalent nucleic acid molecules are referred to as "equivalent nucleic acids." This invention also encompasses nucleic acid molecules characterized by changes in non-coding regions that do not alter the phenotype of the polypeptide produced therefrom when compared to the nucleic acid molecule described above. This invention further encompasses nucleic acid molecules which hybridize to the nucleic acid molecule of the subject invention or its complement. As used herein, the term

"nucleic acid" encompasses mRNA and cRNA as well as single and double-stranded genomic DNA, DNA and cDNA. In addition, as used herein, the term "polypeptide" encompasses any naturally occurring allelic variant thereof, such as S-endoglin and L-endoglin, as well as man-made recombinant forms.

This invention provides an isolated nucleic acid molecule encoding a soluble endoglin-derived polypeptide. As used herein, the term "isolated nucleic acid molecule" means a nucleic acid molecule that is in a form that does not occur in nature. One means of isolating a human endoglin nucleic acid is to probe a human cDNA expression library with a natural or artificially designed antibody to endoglin, using methods well known in the art (see Gougos, A. et al., J. Biol Chem. 265:8361 (1990)) and the Examples set forth below. DNA and cDNA molecules which encode human endoglin polypeptides can be used to obtain complementary genomic DNA, cDNA or RNA from human, mammalian or other animal sources. Isolated genomic DNA also is encompassed by this invention as described above. It is isolated by using the nucleic acid sequences of this invention and methods well known to those of skill in the art as described in Sambrook et al., supra.

The invention further provides an isolated nucleic acid molecule operatively linked to a promoter of RNA transcription, as well as other regulatory sequences. As used herein, the term "operatively linked" means positioned in such a manner that the promoter will direct the transcription of RNA off of the nucleic acid molecule. Examples of such promoters are SP6, T4 and T7. Vectors which contain both a promoter and a cloning site into which an inserted piece of DNA is operatively linked to that promoter are well known in the art. Preferable, these vectors are capable of transcribing RNA in vitro or in vivo. Examples of such vectors are the pGEM series

(Promega Biotec, Madison, WI).

This invention provides an expression or replication vector comprising this isolated nucleic acid molecule such as DNA, cDNA or RNA encoding a soluble endoglin-derived polypeptide. Examples of vectors are viruses, such as bacteriophages, baculoviruses and retroviruses, cosmids, plasmids (such as pcEXV-2) and other recombination vectors. Nucleic acid molecules are inserted into vector genomes by methods well known in the art. For example, insert and vector DNA can both be exposed to a restriction enzyme to create complementary ends on both molecules that base pair with each other and which are then joined together with a ligase. Alternatively, synthetic nucleic acid linkers can be ligated to the insert DNA that correspond to a restriction site in the vector DNA, which is then digested with a restriction enzyme that recognizes a particular nucleotide sequence. Additionally, an oligonucleotide containing a termination codon and an appropriate restriction site can be ligated for insertion into a vector containing, for example, some or all of the following: a selectable marker gene, such as neomycin gene for selection of stable or transient transfectants in mammalian cells; enhancer/promoter sequences from the immediate early gene of human CMV for high levels of transcription; transcription termination and RNA processing signals from SV40 for mRNA stability; SV40 polyoma origins of replication and ColE1 for proper episomal replication; versatile multiple cloning sites; and T7 and SP6 RNA promoters for in vitro transcription of sense and anti-sense RNA. Other means are available.

Also provided are vectors comprising a DNA molecule encoding an endoglin-derived polypeptide, or soluble fragment thereof, adapted for expression in a bacterial cell, a yeast cell, a mammalian cell and other animal cells. The vectors additionally comprise the

regulatory elements necessary for expression of the DNA in the bacterial, yeast, mammalian or animal cells so located relative to the DNA encoding soluble endoglin polypeptide as to permit expression thereof. Regulatory elements  
5 required for expression include promoter sequences to bind RNA polymerase and transcription initiation sequences for ribosome binding. For example, a bacterial expression vector includes a promoter such as the lac promoter and for transcription initiation the Shine-Dalgarno sequence and  
10 the start codon AUG (Sambrook et al. supra. 1989). Similarly, a eucaryotic expression vector includes a heterologous or homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome.  
15 Such vectors can be obtained commercially or assembled by the sequences described in methods well known in the art, for example the methods described above for constructing vectors in general. Expression vectors are useful to produce cells that express the polypeptide.

20 This invention provides a mammalian cell containing a cDNA molecule encoding an endoglin-derived polypeptide or a soluble fragment thereof. An example is a mammalian cell comprising a plasmid adapted for expression in a mammalian cell. The plasmid has a cDNA  
25 molecule encoding an endoglin-derived polypeptide and the regulatory elements necessary for expression of the polypeptide. Various mammalian cells may be utilized as hosts, including, for example, mouse fibroblast cell NIH3T3, CHO cells, HeLa cells, Ltk- cells, etc. Expression  
30 plasmids such as those described supra can be used to transfect mammalian cells by methods well known in the art such as calcium phosphate precipitation, DEAE-dextran, electroporation or microinjection.

This invention provides a pharmaceutical  
35 composition containing a pharmaceutical carrier and any of

a purified polypeptide, a purified soluble polypeptide, an active fragment thereof, or a purified, mature protein and active fragments thereof, alone or in combination with each other. These polypeptides or proteins can be recombinantly  
5 derived, chemically synthesized or purified from native sources. As used herein, the term "pharmaceutically-acceptable carrier" encompasses any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water and emulsions such as an oil/water  
10 or water/oil emulsion, and various types of wetting agents. Any of these pharmaceutical compositions are useful in the methods described below or for the preparation of a medicament for treating the conditions described below.

Also provided are antibodies having specific  
15 reactivity with the endoglin-derived TGF- $\beta$ -binding polypeptides of the subject invention, such as anti-endoglin antibody 44G4 (Quackenbush, E.J., and Letarte, M.J., J. Immunol. 134:1276-1285 (1985)) or any antibody having specific reactivity to a TGF- $\beta$ -binding endoglin  
20 polypeptide. Active fragments of antibodies are encompassed within the definition of "antibody." The antibodies and fragments of the invention can be produced by any method known in the art. For example, polyclonal and monoclonal antibodies can be produced by methods well  
25 known in the art, as described, for example, in Harlow and Lane, Antibodies: A Laboratory Manual (Cold Spring Harbor Laboratory 1988), which is incorporated herein by reference. The polypeptide can be used as the immunogen in generating such antibodies. Altered antibodies, such as  
30 chimeric, humanized, CDR-grafted or bifunctional antibodies can also be produced by methods well known to those skilled in the art. Such antibodies can also be produced by hybridoma, chemical synthesis or recombinant methods described, for example, in Sambrook et al., supra. The  
35 antibodies can be used for determining the presence or purification of the endoglin-derived polypeptide or soluble



fragment thereof, of the present invention. With respect to the detecting of such polypeptides, the antibodies can be used for in vitro diagnostic methods to determine the presence of endoglin or in vivo imaging methods.

5 Immunological procedures useful for in vitro detection of the target soluble endoglin-derived polypeptide in a sample include immunoassays that employ a detectable antibody. Such immunoassays include, for example, ELISA, Pandex microfluorimetric assay,  
10 agglutination assays, radioimmunoassays, flow cytometry, serum diagnostic assays and immunohistochemical staining procedures which are well known in the art. An antibody can be made detectable by various means well known in the art. For example, a detectable marker can be directly or  
15 indirectly attached to the antibody. Useful markers include, for example, radionuclides, enzymes, fluorogens, chromogens and chemiluminescent labels.

This invention provides a method of modifying a biological function mediated by the regulatory activity of  
20 TGF- $\beta$  which comprises contacting a suitable sample containing TGF- $\beta$  with an effective amount of a biologically active endoglin-derived polypeptide, for example soluble endoglin, or a pharmaceutical composition described above.

As used herein, "an effective amount" refers to  
25 an amount of the polypeptide sufficient to bind to TGF- $\beta$  and thereby prevent or inhibit its regulatory activity. This method is especially useful for modifying the regulatory activity of TGF- $\beta$ 1 or TGF- $\beta$ 3. Examples of regulatory activities include, but are not limited to  
30 stimulation of cell proliferation, cell growth inhibition, promotion of extracellular matrix proteins, and regulation of immune functions. TGF- $\beta$  is known to be a potent chemoattractant for monocytes and can induce IL-1, TNF- $\alpha$ , TGF- $\beta$  and surface Fc $\gamma$ RIII, all of which are involved in the

inflammatory response. Conversely, TGF- $\beta$  can deactivate macrophages by inhibiting the antimicrobial activity and the superoxide anion generation, and induce suppression of class II-restricted Ag presentation by macrophages.

- 5           The method can be practiced in vitro or in vivo.  
If the method is practiced in vitro, contacting is effected by incubating the sample with a polypeptide, a protein or a pharmaceutical composition as described above.

- 10           In vitro the novel nucleic acid molecules and antibodies of this invention are useful to detect and quantify the amount of TGF- $\beta$  in a sample isolated from a subject, such as a human patient. The detection of TGF- $\beta$  is useful to monitor the progression of a disease related to overexpression of TGF- $\beta$ , e.g., glomerulonephritis.

- 15           However, in a preferred embodiment the contacting is effected in vivo by administering a polypeptide, a protein or a pharmaceutical composition, as described above, to a subject, e.g., a human patient.

- 20           Methods of administration are well known to those of skill in the art and include, but are not limited to administration orally, intravenously or parenterally. Administration will be in such a dosage such that the regulatory activity is effectively modified. Administration can be effected continuously or  
25           intermittently such that this amount is effective for its intended purpose.

- 30           This invention also provides a method of treating a pathologic condition caused by a TGF- $\beta$ -regulated activity comprising contacting the TGF- $\beta$  with any of a purified soluble endoglin-derived polypeptide, an active fragment thereof, an endoglin-derived polypeptide or an active fragment thereof. The TGF- $\beta$  is bound with said polypeptide

to thereby treat the pathologic condition mediated by TGF- $\beta$  regulatory activity. As used herein, "pathologic conditions" refers to any pathology arising from TGF- $\beta$ -induced regulatory activity, for example, inflammation, rheumatoid arthritis, inflamed skin lesions, scar tissue formation, lung fibrosis, liver fibrosis, atherosclerosis, and glomerulonephritis. Growth and proliferation of mesenchymal cells is stimulated by TGF- $\beta$ , however some tumor cells may also be stimulated thus using TGF- $\beta$  as an autocrine growth factor. An example of inhibitory conditions are the prevention of new cell growth to assist in repair of tissue damage, for ulceration and immunosuppression. The stimulation of extracellular matrix production by TGF- $\beta$  is essential for wound healing. However, in some cases, the TGF- $\beta$  response is uncontrolled and an excessive accumulation of extracellular matrix results. An example of excessive accumulation of extracellular matrix is glomerulonephritis. An additional example of a pathology is cancer.

In one embodiment, the method is practiced by administering to a subject, e.g., a human patient or a mammal, an effective amount of a purified endoglin protein or an endoglin-derived soluble polypeptide or a biologically active fragment thereof, or the pharmaceutical composition described above. Methods of administration are outlined supra.

Also provided by this invention is a method of inhibiting the activity of endoglin by contacting endoglin with an effective amount of a polypeptide capable of binding to endoglin to bind endoglin, thereby inhibiting the activity of endoglin. As used herein, the term "a polypeptide capable of binding to endoglin" means any substance capable of forming a complex with endoglin, for example, TGF- $\beta$ 1 or TGF- $\beta$ 3, or an active fragment thereof. An active fragment is an amino acid sequence corresponding

to a fragment of TGF- $\beta$ 1 or TGF- $\beta$ 3 that retains the ability to bind endoglin. Methods of making such fragments are well known to those of skill in the art as are methods of determining the binding activity of the fragments. Also encompassed by this invention are polypeptides that retain their activity to bind to endoglin, but no longer mediate the biological response corresponding to the binding of a functional ligand to the receptor is destroyed. Thus, these "mutated" polypeptides can act as antagonists to the biological function mediated by the ligand to endoglin by blocking the binding of normal, functioning ligands to endoglin on the cell.

This invention also encompasses the use of the compositions defined above for the preparation of medicaments to modify a biological function regulated by TGF- $\beta$ . These biological functions are described above in detail.

It is understood that modifications which do not substantially affect the activity of the various molecules of this invention are also included within the definition of said molecules.

The following examples are intended to illustrate but not limit the present invention.

**A. ISOLATION OF HUMAN ENDOGLIN PROTEIN  
AND NUCLEIC ACID ENCODING SAME**

**EXAMPLE I**

**Cell Culture and Transfections**

Human umbilical vein endothelial cells (HUVEC, CRL 1730, ATCC) were maintained in  $\alpha$ -minimal essential media supplemented according to supplier's instructions or prepared from umbilical veins as previously described

(Gougos, A. et al., J. Immunol. 141:1925 (1988)). Similar results were obtained using cells from either source. COS-M6 cells, maintained in Dulbecco's modified Eagle's medium supplemented with 10% bovine serum, were transfected with  
5 a cDNA encoding full-length endoglin ligated into the EcoRI site of the mammalian expression vector pcEXV (Miller, J. et al., J. Exp. Med. 164:1478 (1986)) or with a control vector without cDNA insert (pcMV5; Lopez-Casillas, F. et al., Cell 67:785 (1991)) by the DEAE-dextran-chloroquine  
10 procedure (Seed, B., et al., P.N.A.S. USA 84:3365 (1987)). Twenty-four (24) hours post-transfection, cells were trypsinized and reseeded into multicuster dishes and allowed to grow an additional 48 hours before being affinity-labeled with  $^{125}\text{I}$ -TGF- $\beta$ 1 as described below.

15

## EXAMPLE II

Receptor Affinity Labeling and Immunoprecipitation

TGF- $\beta$ 1 and TGF- $\beta$ 2 were purchased from R & D Systems (Minneapolis, MN) and TGF- $\beta$ 3 was obtained from Oncogene Science (Manhasset, NY).  $^{125}\text{I}$ -TGF- $\beta$ 1 used in these  
20 studies was prepared by the chloramine-T method as previously described (Cheifetz, S. et al., J. Biol. Chem. 265:20533 (1990)) or purchased from Amersham Corp.; both preparations gave identical results. The conditions for affinity labeling cell monolayers with  $^{125}\text{I}$ -TGF- $\beta$ 1 and  
25 disuccinimidyl suberate (Pierce Chemical Co.) have been described previously (Massague, J., Methods Enzymol. 146:174 (1987)). The concentrations of  $^{125}\text{I}$ -TGF- $\beta$ 1 and competing unlabeled ligands used for each experiment are indicted in the figure legends. Triton X-100 extracts of  
30 the affinity-labeled cells were either analyzed directly on sodium dodecyl sulfate-polyacrylamide gels (SDS-PAGE) or first incubated with monoclonal antibody (mAb) 44G4 directed against human endoglin (Quackenbush, E.J. et al., J. Immunol. 134:1276 (1985)) or with control antibody (see  
35 below). For immunoprecipitations, detergent extracts were

diluted with an equal volume of phosphate-buffered saline containing 1% Triton X-100 and precleared by incubation for 20 min at 4°C with protein G-Sepharose (Pharmacia LKB Biotechnology Inc.) prior to overnight incubation at 4° C. with mAb 44G4. Immune complexes were collected by incubation with protein G-Sepharose for 1 hour at 4° C. For some experiments, mAb 44G4 was used coupled to Sepharose. The immunoprecipitates were washed three times (saline with 1% Triton X-100) and then resolved by SDS-PAGE in the presence or absence of dithiothreitol (DTT) and visualized by autoradiography. Irrelevant mAb (44D7) used in control experiments to monitor specificity of the immunoprecipitations did not immunoprecipitate any affinity-labeled bands.

15

## EXAMPLE III

SDS-PAGE and 2D-Gel Analysis

Analysis of the affinity-labeled profile of HUVEC revealed that, like vascular endothelial cells from other sources, these cells have little or no betaglycan, which characteristically migrates as a diffuse band between 200 and 400 KDa on reducing SDS-PAGE (Figure 2A). Instead, HUVEC expressed a disulfide-linked cell surface protein that, together with TGF- $\beta$  receptors I and II, was affinity-labeled by crosslinking with  $^{125}$ I-TGF- $\beta$ 1. Receptors I and II were detected in HUVEC as labeled complexes of approximately 65 KDa and 100 KDa, respectively, which is similar to the size of these labeled receptors reported for other human cell lines. Comparison of the relative migration of the affinity-labeled proteins fractionated on SDS-PAGE revealed that the major affinity-labeled proteins of HUVEC migrated between 95-120 KDa on reducing gels whereas on non-reducing gels the major affinity-labeled proteins migrated between 100-110 KDa (presumed to be receptor II) and at 180 KDa and above (endoglin) (Figure 2). This pattern indicated the presence of disulfide-

linked TGF- $\beta$ -binding proteins.

Resolution of these disulfide-linked TGF- $\beta$ 1 binding proteins on two-dimensional gels (Figure 2B) confirmed that the disulfide-linked complexes (probably dimers and higher order oligomers) contained subunits of approximately 95 KDa (value estimated by subtracting the cross-linked TGF- $\beta$ 1 monomer mass 12.5 KDa from the reduced 110 KDa affinity-labeled complex). Together with the type II receptor, the disulfide-linked TGF- $\beta$ 1-binding proteins are the major affinity-labeled species expressed by HUVEC.

#### EXAMPLE IV

##### Immunoprecipitation with anti-endoglin mAb

To determine whether the disulfide-linked TGF- $\beta$ -binding protein on endothelial cells was endoglin, affinity-labeled HUVEC extracts were immunoprecipitated with monoclonal antibody (mAb) 44G4 which is specific for human endoglin (Georgi, L.L. et al., Cell 61:635 (1990); MacKay, K. et al., J. Biol. Chem. 266:9907 (1992); Merwin, J.R. et al., Am. J. Pathol. 138:37 (1991)). Electrophoretic analysis of these immunoprecipitates revealed a labeled protein complex whose subunit structure was similar to that of endoglin (Figure 3A). Thus, under reducing conditions, a major affinity-labeled band of approximately 110 KDa was seen which migrated as complexes of 180 KDa and greater than 200 KDa when analyzed under non-reducing conditions. The higher order oligomers might contain multiple endoglin molecules crosslinked by TGF- $\beta$ 1, itself a disulfide-linked dimer. Repeated immunoprecipitation with 44G4-IgG-Sepharose completely depleted these labeled species from cell extracts (Figure 3B). No affinity-labeled bands were immunoprecipitated from three other human cell lines (A549, Hep G2, MCF-7), which lack endoglin and were used as negative-controls for these experiments. Monoclonal antibodies specific to human

endoglin thus demonstrate that endoglin is a major TGF- $\beta$ -binding protein in human vascular endothelial cells.

#### EXAMPLE V

##### Ectopic Expression of Endoglin in Cells

5           The identity of this dimeric TGF- $\beta$ -binding protein of HUVEC with endoglin was confirmed by ectopically expressing the full-length endoglin cDNA in COS monkey kidney cells. After affinity-labeling with  $^{125}\text{I}$ -TGF- $\beta$ 1, a labeled species with the characteristics of endoglin could  
10 be specifically precipitated by mAb 44G4 only from the detergent extracts of endoglin transfectants (Figure 4). Differences in glycosylation likely account for the smaller size of endoglin expressed in COS cells relative to endogenous endoglin of HUVEC.

#### 15           B. ISOLATION OF S- AND L-ENDOGLIN

#### EXAMPLE VI

##### cDNA Cloning and Sequencing of Endoglin

Approximately  $2.5 \times 10^5$  clones from a  $\lambda$ gt10 cDNA library prepared from PMA-treated myelomonocytic human cell  
20 line HL60 (Corbi, A.L. et al., EMBO J. 6:4023 (1987)). (50,000 pfu/150 mm dish) were screened with 700 bp PstI fragment from endoglin cDNA (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)). Thirteen hybridizing clones were isolated after three rounds of plaque  
25 purification with sizes ranging from 2.4-3.1 kb. The longest clone (clone 3.3; 3073 bp) was subcloned into pUC13 and sequenced using the dideoxy chain termination method (Sanger, F. et al., PNAS USA 74:5463 (1977)).

Clone 3.3 in pUC13 was digested with BbrPI and  
30 BamHI. Endoglin fragment was made blunt and inserted into the mammalian expression vector pcEXV (Miller, J. and



Germain, R., J. Exp. Med. 164:1478 (1986)), yielding pcEXV-EndoS. The lack of leader sequence in the cDNA (Table I) (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990) was overcome by the construction of pcEXV-EndoL.

5 pcEXV-EndoS was digested with MluI/BamHI and ligated to the 563 bp MluI/BamHI fragment specific to endoglin cDNA (Table I), resulting in pcEXV-EndoL. Transfectants were generated by cotransfection of either pcEXV-EndoS or pcEXV-EndoL with psV2neo into mouse L cells. After G418 selection,

10 endoglin-positive clones were isolated. In the same experiment, endoglin-negative clones were selected as mock transfectants.

#### EXAMPLE VII

##### 15 Antibodies, Immunofluorescence, Immunoprecipitations and Immunoblotting

Endoglin-specific monoclonal antibodies used were 8E11 (Lastres, P. et al., Eur. J. Immunol. 22:393 (1992)) of the IgM class and 44G4 (Gougos, A. and Letarte, M., J. Immunol. 141:1925 (1988)) of the IgG1 subclass. Control

20 monoclonal antibodies were HC1/1 (anti-CD11c) of the IgG1 subclass and X63. FCM and immunoprecipitation analyses were performed as described in Lastres, P. et al., Eur. J. Immunol. 22:393 (1992), incorporated herein by reference. For immunoblotting studies,  $5 \times 10^6$  cells were lysed in 250

25  $\mu$ l of 1% Triton X-100, 1 mM PMSF in PBS for 30 minutes at 4°C. Insoluble material was removed by centrifugation at 100,000 x g for 1 hour in a Beckman TL100 centrifuge. Proteins contained in the supernatant were separated by

30 SDS-PAGE under nonreducing conditions using a minigel system and transferred to nitrocellulose membranes (Hybond, Amersham). Membranes were incubated first with a blocking solution (10% fetal calf serum, 0.5% Tween-20, 1 M glucose and 10% glycerol in PBS), followed by incubation with the monoclonal antibody 44G4. The presence of endoglin was

35 revealed using a chemiluminescence assay (ECL detection

kit, Amersham).

#### EXAMPLE VIII

##### Reverse Transcription-Polymerase Chain Reaction (RT-PCR)

Total cellular RNA was isolated using guanidinium isothiocyanate and a cesium chloride ultracentrifugation step (Chirgwin, J.M. et al., Biochemistry 18:5294 (1979)). Poly (A)+ RNA was purified by oligo (dT) affinity chromatography (Maniatis, T. et al., "Molecular Cloning: A Laboratory Manual" Cold Spring Harbor Laboratory, Cold Spring Harbor (1982). Single-stranded cDNA was synthesized from poly (A+) RNA (0.5 µg/40 µl reaction), using AMV reverse transcriptase. Five microliters of cDNA was used for a 50 µl PCR reaction (Gilliland, G. et al., In: [Innis, M.A., Gelfand, D.H., Sninsky, J.J. and White, T.J. (Eds.)] "PCR Protocols" Academic Press, San Diego 1990, p. 60.). Oligonucleotide primers used for amplification were E#12 (nucleotides 1945-1964), E#11R (reverse complement of nucleotides 2475-2494), E#14 (nucleotides 2136-2158), and E#15 (reverse complement of nucleotides 2247-2273). Amplifications were performed in 1 X Taq buffer (Promega) with 0.2 mM each dATP, dCTP, dGTP, dTTP at 0.25 µM concentration of each primer and 0.25 U/50 µl of Taq DNA polymerase (Promega). Amplification was carried out in a thermal cycler as follows: 5 minutes at 95°C; 35 cycles of 45 seconds at 94°C, 45 seconds at 54°C, and 1 minute at 72°C for oligonucleotide pair E#12-E#11R or 35 cycles of 45 seconds at 94°C, 45 seconds at 68°C and 1 minute at 72°C for pair E#14-E#15 and then 10 minutes at 72°C. Control reactions, identical except for the omission of reverse transcriptase, were performed simultaneously. Amplification products were analyzed by agarose gel electrophoresis and ethidium bromide staining.

## EXAMPLE IX

Receptor Affinity Labeling

TGF- $\beta$ 1 was purchased from R&D Systems (Minneapolis, MN) and  $^{125}$ I-TGF- $\beta$ 1 was obtained from Amersham (Oakville, Canada) at a specific activity of 2000 Ci/mmol. Parental mouse L cells, transfectant mouse L cells expressing S-endoglin or L-endoglin, and mock transfectant L cells were grown at confluence ( $5 \times 10^6$  cells per plate) and incubated with 100 pM of  $^{125}$ I-TGF- $\beta$ 1 for 4 hours with and without 4 nM of competing unlabeled TGF- $\beta$ 1. Cells were washed and cross-linked at a final concentration of 0.16 mM disuccinimidyl suberate (DSS) (Pierce Chemical Co.) in a buffer containing 128 mM NaCl, 5 mM KCl, 5 mM MgSO<sub>4</sub>, 1.2 mM CaCl<sub>2</sub>, 50 mM Hepes, pH 7.5 for 15 minutes at 4°C. Cells were washed four times and solubilized directly on the petri dishes with a minimum of solubilization buffer containing 1% Triton X-100 and proteolytic inhibitors as described previously (Massagué, J., Methods Enzymol. 146:174 (1987)). The extracts were subjected to immunoprecipitation followed by SDS-PAGE analysis.

## RESULTS

Isolation and Characterization of Endoglin cDNA Clones:Identification of a Cytoplasmic Variant

Full-length cDNA clones were derived from a  $\lambda$ gt10 library prepared from PMA-treated HL60 cells. Clone number 3.3 was selected for further characterization due to the large size of the insert. The partial sequence of the 3073 bp cDNA insert from clone 3.3 is depicted in Figure 7. This clone contains 281 bp of 5' untranslated region followed by an open reading frame of 1875 bp. The predicted protein sequence shows that the endoglin leader peptide contains 25 amino acids (aa), followed by 561 residues at the extracellular portion and a transmembrane

region spanning 25 amino acids as expected (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)). However, clone 3.3 contains a 135 bp segment inserted within the previously known cDNA sequence, starting at nucleotide (nt) 2134 (Figure 8). The first 21 nucleotides of the insert are in frame with the preceding sequence and code for a new sequence of 7 amino acids. This sequence replaces the C-terminal 40 amino acids of endoglin, thus leading to a new cytoplasmic tail of 14 amino acids different from the 47 residues previously reported (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)) (Figure 8), suggesting the existence of two alternative forms of endoglin. The predominant form of endoglin in the myelomonocytic cell line HL60 seems to be the one containing 47 residues in the cytoplasmic domain since this sequence was present in 12 out of 13 clones analyzed. The isoforms with the 14 and 47 amino acid cytoplasmic tails, and the corresponding cDNAs, are referred to as S-endoglin and L-endoglin, respectively.

#### Expression of two different forms of endoglin

To characterize further the two alternative forms, independent constructs corresponding to the short and long forms of endoglin were inserted into the expression vector pcEXV and transfected into mouse fibroblasts (Figure 9). Both, S-endoglin and L-endoglin were highly expressed on the cell surface as determined by FCM (Figure 9). In addition, metabolic labeling of the transfectants followed by immunoprecipitation, revealed a 170-kDa (L-endoglin) or a 160-kDa (S-endoglin) protein specifically recognized by anti-endoglin monoclonal antibody (Figure 9). The distinct size of S- and L-endoglin in the transfectants was also detected by immunoblotting analysis under non-reducing conditions (Figure 9). In these experiments, L-endoglin showed the same  $M_r$  as the endoglin detected on U937 cells, suggesting that this is the predominant form in the promonocytic cell

line.

Cell surface radiolabeling of the transfectants followed by immunoprecipitation demonstrated that both isoforms are expressed as disulfide linked homodimers (Figure 9), indicating that cysteine residue present in the extracellular portion are mediating the interchain disulfide bond.

Differential expression  
of S-endoglin and L-endoglin mRNA

10           The individual expression of S- and L-endoglin  
mRNA was analyzed by RT-PCR. Amplification with two  
primers derived from the unique 3' region of S-endoglin  
cDNA generated the expected 137 bp product on S-endoglin  
transfectants and on PMA-treated monocytic lines,  
15   endothelial cells and placenta (Figure 10), demonstrating  
the existence of this isoform in several cell types. When  
primers common to L-endoglin and S-endoglin were used, the  
L-endoglin specific fragment of 411 bp could be amplified  
on PMA-treated monocytic cell lines, placenta and  
20   endothelial cells, whereas the S-endoglin fragment of 546  
bp was only amplified on control S-endoglin transfectants  
(Figure 10). When competitive templates are amplified by  
PCR, an abundant template can suppress the amplification of  
a less abundant one (Gilliland, G. et al., In: [Innis,  
25   M.A., Gelfand, D.H., Sninsky, J.J. and White, T.J. (Eds.)  
"PCR Protocols", Academic Press, San Diego 1990, p. 60].  
Therefore, these results indicate that L-endoglin is the  
predominant form and S-endoglin is expressed at lower  
levels in these cell types. This is in agreement with the  
30   presence in PMA-treated U937 cells of an endoglin molecule  
of  $M_r$  similar to that observed on the transfectant L-  
endoglin (Figure 9), assuming similar glycosylation levels  
on both cell types.

L-endoglin and S-endoglin bind to TGF- $\beta$ 1

Since endoglin has been found to be a component of the receptor system for TGF- $\beta$ , Cheifetz, S. et al., J. Biol. Chem. 267:19027 (1992), it was of interest to analyze the ability of each isoform to bind this ligand. Transfectants expressing short and long forms of endoglin were compared to the parental L cells and to the mock transfectants for their ability to bind TGF- $\beta$ 1. Immunoprecipitation analysis allowed the identification of the TGF- $\beta$ 1-endoglin complex (Figure 11). Under non-reducing conditions, the dimers of endoglin were seen as radiolabelled bands of  $M_r$  170 kDa (short form) and 175 kDa (long form). An oligomer of endoglin is seen in both cases migrating with a  $M_r$  > 270 kDa. A similar complex was also observed previously in cross-linking experiments with human endothelial cells and might represent endoglin molecules cross-linked by TGF- $\beta$ 1, itself a dimer (Roberts, A.B. and Sporn, M.B. In: Sporn, M.B. and Roberts, A.B. (Eds.), "Peptide growth factors and their receptors" Springer-Verlag, Heidelberg 1990, p. 419). Under reducing conditions, major bands of 97 kDa and 107 kDa were seen with the short form of endoglin and bands of 102 kDa and 112 kDa were observed with the long form. These doublets have been observed previously for endothelial cells and might represent endoglin bound to the monomer (12.5 kDa) or dimer (25 kDa) of TGF- $\beta$ 1 (Roberts, A.B. and Sporn, M.B. In: Sporn, M.B. and Roberts, A.B. (Eds.), "Peptide growth factors and their receptors" Springer-Verlag, Heidelberg 1990, p. 419.; Cheifetz, S. and Massagué, J., J. Biol. Chem. 266:20767 (1991)). By subtracting the contribution of TGF- $\beta$ 1, one can estimate a molecular weight of 85 kDa for S-endoglin and 90 kDa for L-endoglin.

DISCUSSION

Sequence analysis demonstrated the existence of two different cDNA variants named L-endoglin and S-endoglin. These two isoforms are coexpressed by myeloid cells, endothelium and placenta, although the majority of the transcripts synthesized apparently correspond to the L-endoglin isoform. The mechanism by which the two isoforms are generated remains to be determined. Most likely, both isoforms are generated by alternative splicing. In fact, consensus sequences of donor/acceptor sites (GT, AG) at the 5' and 3' ends and branch point of the lariat (CTGAC), have been found on the novel 135 bp insert of S-endoglin cDNA (Figure 7). A similar example of cytoplasmic variants generated by a "retained intron" splicing mechanism, have been recently reported for the activin receptor of the TGF- $\beta$  receptor family (Attisano, L. et al., Cell 68:97 (1992)).

The behavior of the two endoglin isoforms was analyzed by transfection studies. Both forms are expressed on the cell surface as disulfide linked homodimers, indicating that the cysteine residues present in the extracellular region are responsible for the dimerization. In spite of the different cytoplasmic domains, both forms behave as TGF- $\beta$  binding proteins.

S-endoglin, L-endoglin and betaglycan contain 22 identical residues in the transmembrane domain and the adjacent cytoplasmic region (Figure 9). This delineates a first conserved motif between betaglycan and the two forms of endoglin, which contains two tyrosine residues and two Ser/Thr residues whose phosphorylation status remain to be analyzed. A second region of high identity in the cytoplasmic tail is shared only by the long form of endoglin and betaglycan. The high content (40%) of Ser/Thr residues in this motif, and the absence of Tyr suggests that this region might undergo phosphorylation by a Ser/Thr

kinase. Interestingly, the cytoplasmic domain of the TGF- $\beta$  receptor II, displays Ser/Thr kinase activity (Lin, H.Y. et al., Cell 68:775 (1992)).

Although the invention has been described with  
5 reference to the disclosed embodiments, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.



What is claimed is:

1. An isolated nucleic acid molecule encoding an endoglin polypeptide or a biologically active fragment thereof.
- 5 2. A nucleic acid molecule encoding a soluble endoglin-derived polypeptide or a biologically active fragment thereof.
3. The nucleic acid of claim 1 or 2, wherein the nucleic acid encodes a human endoglin polypeptide.
- 10 4. The nucleic acid molecule of claim 1 or 2, wherein the nucleic acid is genomic DNA, cDNA, mRNA or cRNA.
5. The nucleic acid molecule of claim 4, wherein the genomic DNA is designated CECT 4475.
- 15 6. The nucleic acid molecule of claim 1 or 2 operatively linked to a promoter of RNA transcription.
7. A vector containing the nucleic acid molecule of claim 5.
8. A host cell containing the vector of claim
- 20 7.
9. The host cell of claim 8, wherein the cell is a procaryotic cell or a eucaryotic cell.

10. A method of preparing an endoglin-derived polypeptide or active fragment thereof, comprising:

- a. inserting a nucleic acid molecule encoding an endoglin-derived polypeptide or active fragment thereof  
5 into a suitable expression vector;
  - b. inserting the resulting vector into a suitable host cell;
  - c. inducing the resulting host cell to express the endoglin-derived polypeptide or active fragment  
10 thereof; and
  - d. purifying the resulting endoglin-derived polypeptide so produced.
11. Soluble endoglin-derived polypeptide.
12. Endoglin-derived polypeptide or a  
15 biologically active fragment thereof produced by the method of claim 10.
13. A pharmaceutical composition comprising the polypeptide of claim 11 or 12 and a pharmaceutically acceptable carrier.
- 20 14. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a purified and isolated human endoglin-derived polypeptide or a biologically active fragment thereof substantially free of other host cell proteins.

15. A method of modifying a biological function mediated by the regulatory activity of TGF- $\beta$  which comprises contacting TGF- $\beta$  with an effective amount of an endoglin-derived polypeptide or a biologically active  
5 fragment thereof, thereby modifying the biological function.

16. The method of claim 15, wherein TGF- $\beta$  is TGF- $\beta$ 1.

17. The method of claim 15, wherein TGF- $\beta$  is  
10 TGF- $\beta$ 3.

18. The method of claim 15, wherein the contacting is effected in vitro.

19. The method of claim 15, wherein the contacting is effected in vivo.

20. The method of claim 15, wherein the regulatory activity is stimulation of cell proliferation or cell growth inhibition.

21. The method of claim 15, wherein the regulatory activity is promotion of extracellular matrix  
20 production.

22. The method of claim 15, wherein the regulatory activity is regulation of immune function.

23. The method of claim 15, wherein the polypeptide is a purified human endoglin-derived  
25 polypeptide comprising a disulfide-linked homodimer of about 80 to about 95 kDa subunits or a biologically active fragment thereof.

24. The method of claim 23, wherein the purified polypeptide is purified and isolated human endoglin-derived protein substantially free of other human proteins.

25. The method of claim 15, wherein the polypeptide is a soluble endoglin-derived polypeptide.

26. A method of treating a pathologic condition caused by TGF- $\beta$  regulated cell growth stimulation which comprises administering to a subject an effective amount of the pharmaceutical composition of claim 13 or 14 to bind to the TGF- $\beta$  thereby treating the pathologic condition caused by cell growth stimulation.

27. The method of claim 26, wherein TGF- $\beta$  is TGF- $\beta$ 1.

28. The method of claim 26, wherein TGF- $\beta$  is TGF- $\beta$ 3.

29. The method of claim 26, wherein the subject is a human patient.

30. A method of treating a pathologic condition caused by TGF- $\beta$  regulated inhibition of cell growth which comprises administering to a subject an effective amount of the pharmaceutical composition of claim 13 or 14 to bind to the TGF- $\beta$ , thereby treating the pathologic condition caused by inhibition of cell growth.

31. The method of claim 30, wherein TGF- $\beta$  is TGF- $\beta$ 1.

32. The method of claim 30, wherein TGF- $\beta$  is TGF- $\beta$ 3.

33. The method of claim 30, wherein the pathologic condition is ulceration or immune suppression.

34. The method of claim 30, wherein the subject is a human patient.

5           35. The method of treating a pathologic condition caused by TGF- $\beta$  regulated promotion of extracellular matrix accumulation which comprises administering to a subject an effective amount of the pharmaceutical composition of claim 12 or 13 to bind TGF- $\beta$   
10 thereby treating the pathologic condition caused by promotion of extracellular matrix.

36. The method of claim 35, wherein the subject is a human patient.

15           37. The method of claim 35, wherein the pathologic condition is inflammation, rheumatoid arthritis, inflamed skin lesions, scar tissue formation, lung fibrosis, liver fibrosis, atherosclerosis or glomerulonephritis.

20           38. A method of inhibiting the activity of endoglin which comprises contacting endoglin with an effective amount of a polypeptide capable of binding to endoglin to bind endoglin, thereby inhibiting the activity of endoglin.

25           39. The method of claim 38, wherein the polypeptide is a TGF- $\beta$ .

40. The method of claim 38, wherein the polypeptide has an amino acid sequence corresponding to a fragment of a TGF- $\beta$  having the ability to bind endoglin.

41. The method of claim 39, wherein the TGF- $\beta$  is TGF- $\beta$ 1 or TGF- $\beta$ 3.

42. The method of claim 40, wherein the TGF- $\beta$  is TGF- $\beta$ 1 or TGF- $\beta$ 3.

5 43. The method of claim 38, wherein the polypeptide is an anti-endoglin antibody.

44. The method of claim 43, wherein the antibody is a monoclonal antibody.

10 45. The method of claim 38, wherein the contacting is effected in vivo.

46. The method of claim 38, wherein the contacting is effected in vitro.

- 1 / 13

FIGURE 1






FIGURE 2





- 3 / 13

FIGURE 3



- 4 / 13

FIGURE 4






FIGURE 5



**pcNeoSolEND**  
**8.7 Kb**



**Vector pcDNA1/Neo**  
**7.0 Kb**

[illegible]

FIGURE 8




FIGURE 9







FIGURE 10






FIGURE 11





FIGURE 12



1 : TGGGCGCCAGGACTGCTGCTGTCACTGCCATCCATTGGAGCCGACGACCCCTCCCGCCCATCTTCGGACAGCAACTCCAGCCGACCCG  
91 : CCGCTGCTGTGTTCACCTTCTCTGACCCCTTGGCCGCCACCCAGAGGCTGGAGCAGGGACCGCGTGGCTCCGGCCGCTGCTCTCT  
181 : CCGGCTCCCGCTGGAGCCACGCGCCGCGCGCTGCCCGCCGAGCCCTGCCACTGGACACAGGATTAAGCCCGCCAGCCGAGCCCAAC  
271 : GTGGACAGCATGGACCCCGCCACGCTCCCTCTGGCTGTGGCTGTGGCCAGTTCGAGCCCAAGTCTTCGAGAAACA  
: .....MetAspArgGlyThrLeuProLeuAlaValAlaLeuLeuAlaSerCysSerLeuSerProThrSerLeuAlaGluThr  
361 : GTCCATTGTGACCTTCAGCCTGTGGGCCCCGAGAGGGGGCAGAGGTGACATATACCACTAGCCAGGCTTCGAAGGGCTGGCTGGCTCAGGCC  
28 : ValHisCysAspLeuGlnProValGlyProGluArgGlyGluValThrTyrThrThrSerGlnValSerLysGlyCysValAlaGlnAla  
451 : CCAATGCCATCTCTGAAGTTCATGTCTCTTCTTGGAGTTCACACGGGCGCTCACAGCTGGAGCTGACTTCTCCAGGCATCCAGCAA  
59 : ProAsnAlaIleLeuGluValHisValLeuLeuGluPheProThrGlyProSerGlnLeuGluLeuThrLeuGlnAlaSerLysGln  
541 : AATGGCACTTGGCCCCGAGAGGTGCTTCTGGTCTCAGTGTAACAGCAGTGTCTTCTGCACTCCAGGCCCTGGGAATCTCCACTGGAC  
98 : AsnGlyThrTrpProArgGluValLeuLeuValLeuSerValAsnSerSerValPheLeuHisLeuGlnAlaLeuGlyProLeuHis  
631 : TTGGCTACAATTTCAGCCTGGTACCTTCCAAGAGCCCCGGGGTCAACACCACAGAGCTGCCATCTTCCCCAAGACCAGATCTCTT  
118 : LeuAlaTyrAsnSerSerLeuValThrPheGlnGluProProGlyValAsnThrThrGluLeuProSerPheProLysThrGlnIleLeu  
721 : GAGTGGCGAGCTGAGAGGGGGCCCCATCACCCTCTGCTGCTGAGCTGAATGACCCCCAGAGCATCTCTCTCCGACTGGGCCAAGCCAGGGG  
148 : GluTrpAlaAlaGluArgGlyProIleThrSerAlaAlaGluLeuAsnAspProGlnSerIleLeuLeuArgLeuGlyGlnAlaGlnGly  
811 : TCATGTCTCTTCGATGCTGGAAGCCAGGAGCATGGGCGGACGCTCGAGTGGCGGCGCGGTACTCCAGCCTTGGTCCGGGGCTGC  
178 : SerLeuSerPheCysMetLeuGluAlaSerGlnAspMetGlyArgThrGluLeuTrpArgProArgThrProAlaLeuValArgGlyCys  
301 : CACTTGAAGGCGTGGCCGGCCACAGGAGGCGGCACATCTGAGGGTCTTCCGGGCCATCTCGGCGGGCCCCGGAGCGGTGACGGTGAAG  
208 : HisLeuGluGlyValAlaGlyHisLysGluAlaHisIleLeuArgValLeuLeuGlyHisSerAlaGlyProArgThrValThrValLys  
991 : GTGGAAGTGAAGTGGCAGCCCGGGATCTCSATGCCGTCTCATCTCTGAGGCTCCCGCTACGTGTCTGGCTCATCGAGCCCAACCC  
238 : ValGluLeuSerCysAlaProGlyAspLeuAspAlaValLeuIleLeuGlnGlyProProTyrValSerTrpLeuIleAspAlaSerHis  
1081 : AACATGCAGATCTGGACCACTGAGAAATACTCTTCAAGATCTTTCCAGAGAAAACATTCTGGCTTCAAGCTCCCGAGACACACTCAA  
268 : AsnMetGlnIleTrpThrThrGlyGluTyrSerPheLysIlePheProGluLysAsnIleArgGlyPheLysLeuProAspThrProGln  
1171 : GGCTCTCTGGGGAGGCGGGATGCTCAATGCCAGCATTGTGGCATCTCTCTGGAGCTACCGCTGGCCAGCATGTGCTCACTTCATGCC  
298 : GlyLeuLeuGlyGluAlaArgMetLeuAsnAlaSerIleValAlaSerPheValGluLeuProLeuAlaSerIleValIleSerLeuHisAla  
1261 : TCCAGCTCGCGTGGTAGGCTCCAGACCTCACCCGACCCAGTCCAGACCACTCTTCCCAAGGACACTTGTAGCCCGGAGCTGCTCATGTCC  
328 : SerSerCysGlyGlyArgLeuGlnThrSerProAlaProIleGlnThrThrProLysArgThrCysSerProGluLeuLeuMetSer  
1351 : TTGATCCAGACAAAGTGTCCGACGAGCCATGACCTGGTACTAAGAAAGAGCTTGTTCGCAATTGTGAAGTCACCATCAGGGCGCTG  
358 : LeuIleGlnThrLysCysAlaAspAspAlaMetThrLeuValLeuLysLysGluLeuValAlaHisLeuLysCysThrIleThrGlyLeu  
1441 : ACCTTCTGGGACCCAGCTGTGAGGCAGAGGACAGGGGTGACAAGTTTGTCTTGGCAGTGCTTACTCCAGCTGTGGCATGCAGGTGTCA  
388 : ThrPheTrpAspProSerCysGluAlaGluAspArgGlyAspLysPheValLeuArgSerAlaTyrSerSerCysGlyMetGlnValSer  
1531 : GCAAGTATGCATCAAGGCGGGTGGTCAATCTCTCGAGCTCATCACACAGGGGAAAAGGTGCACGTGCTTCAACATGGACAGC  
418 : AlaSerMetIleSerAsnGluAlaValValAlaSerSerSerSerProGlnArgLysLysValHisCysLeuAsnMetAspSer  
1621 : CTCTCTTTCAGCTGGGCTCTACCTCAGCCACACTCTCTCCAGGCTTCCACACCATTCAGGCGGGGACAGAGCTTGTGTCAGGCT  
448 : LeuSerPheGlnLeuGlyLeuTyrLeuSerProHisPheLeuGlnAlaSerAsnThrIleGluProGlyGlnGlnSerPheValGlnVal  
1711 : AGAGTGTCCCATCCGTCTCCGAGTCTCTGCTCCAGTTAGACAGCTGCCACCTGGACTTGGGGCTGAGGGAGGCACCGTGGAACTCATC  
478 : ArgValSerProSerValSerGluPheLeuLeuGlnLeuAspSerCysHisLeuAspLeuGlyProGluGlyGlyThrValGluLeuIle  
1801 : CAGGGCGGGGCGGCCAAGGGCAACTGTGTAGCCTGCTGTCCCCAAGCCCGAGGGTGACCCGCGCTTACGTTCTCTCTCCACTTCTAC  
508 : GlnGlyArgAlaAlaLysGlyAsnCysValSerLeuLeuSerProSerProGluGlyAspProArgPheSerPheLeuLeuHisPheThr  
1891 : ACAGTACCATACCCAAACCGGCACTTCAGCTGCACGGTAGCCCTCGCTCCCAAGACCGGGTCTCAAGACCAGGAAGTCCATAGGACT  
538 : ThrValIleProIleProAlaThrGlyThrLeuSerCysThrValAlaLeuArgProLysThrGlySerGlnAspGlnGluValHisArgThr  
1981 : GTCTCATGCGCTTGAACATCATCAGCCCTGACCTGTCTGGTTGCACAAGCAAGGCTCTGCTGCCCGCGGTGCTGGGCACTCACTT  
568 : ValPheMetArgLeuAsnIleIleSerProAspLeuSerGlyCysThrSerLysGlyLeuValLeuProAlaValGluGlyIleThrPhe  
2071 : GGTGCTTCTCATCGGGCCCTGCTCACTGCTGCACCTTGGTACATCTACTCGCACAGCGTTCGCCAGCAAGCGGGAGCCGCTGGTG  
598 : GlyAlaPheLeuIleGlyAlaLeuLeuThrAlaAlaLeuTrpTyrIleTyrSerHisThrArgSerProSerLysArgGluProValVal  
2161 : CGGTGGCTGCCCGGCTCTCTCGAGAGCAGCAGCACCACACAGCATCGGAGCAGCCAGAGCAGCCCTGCTCCACCAGCAGCATG  
628 : AlaValAlaAlaProAlaSerSerGluSerSerSerThrAsnHisSerIleGlySerThrGlnSerThrProCysSerThrSerSerMet  
2251 : GCATAGCCCCCGCCCCCGGCTCGCCGACGAGGAGACTGAGCAGCCCGCAGCTGGGAGCACTGGTGTGAACCTACCCCTGGGAGCCAG  
658 : AlaStop.....  
2341 : TCTCTCACTCGACCCAGAAATGGAGCGTGCTCTTCCCGGCTTACCTTCCCGGCTGCTTCTCAGAGGCTCTGCTGCCAGTCCAGCCACTGCGCT  
2431 : TGGACACACTTGGGGTCTCTCCACCCACAGAACTTCAACCACTGGGTCTGGGATATGCTCTCCAGAGGACAGACCACTTGGCAGCC  
2521 : TGTGTAAACAACTGCTCTGCTATTTGAACCTGGATCCAGCACTGGTGAACCTGAGCTGGGACAGGAGGCGCCACAGCACTTGAACAAATT  
2611 : AGCCGACCCAGCCAGGACCCACCTGACCTCTCCCGCTGAGGAGAGAGGCGCCACAGGACCACTTGAATCTATCTCTCGGGCTCTC  
2701 : ACATCTGAACCTGCTTAACTAACTGCGCAGCGGACAGGACCTGAGGAGCTAGCCGAGCTAGCTTGGGAGCAGAGGTTGAGAACAGTGGGG  
2791 : TCTGGAGCACTAGCTCTGCCACATGCGGCCCTCTCGCGCTGGGAGCAGCAGAGGGGGAGTACGCAAGCTGCTTGTCTGGGCT  
2881 : GCGCTGTGTATTCAACACCAATAATTCAGACCATTAACCACTGAAACAGTGAACAAAAA

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/10307

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(5) : Please See Extra Sheet. US CL : 435/70.1, 240.2, 252.3, 320.1; 514/2; 530/395; 536/23.5 According to International Patent Classification (IPC) or to both national classification and IPC																				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/70.1, 240.2, 252.3, 320.1; 514/2; 530/395; 536/23.5 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, Medline, Biosis search terms: endoglin, TGF-B																				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>																				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																		
X — Y — A	J. Biol. Chem., Vol. 265, Number 15, issued 25 May 1990, Gougos et al., "Primary structure of endoglin, an RGD-containing glycoprotein of human endothelial cells", pages 8361-8364, see Fig. 4.	1-4 — 5-12 — 13-46																		
X — A	J. Biol. Chem., Vol. 267, Number 27, issued 25 September 1992, Cheifetz et al., "Endoglin is a component of the transforming growth factor-Beta receptor system in human endothelial cells", pages 19027-19030, see Fig. 4.	6, 10-12 — 13-46																		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.																				
<table border="0"> <tr> <td>* Special categories of cited documents:</td> <td>"T"</td> <td>later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"A" document defining the general state of the art which is not considered to be part of particular relevance</td> <td>"X"</td> <td>document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"E" earlier document published on or after the international filing date</td> <td>"Y"</td> <td>document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)</td> <td>"A"</td> <td>document member of the same patent family</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td></td> <td></td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> <td></td> </tr> </table>			* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"A" document defining the general state of the art which is not considered to be part of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"E" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	"A"	document member of the same patent family	"O" document referring to an oral disclosure, use, exhibition or other means			"P" document published prior to the international filing date but later than the priority date claimed		
* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																		
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																		
"E" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																		
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	"A"	document member of the same patent family																		
"O" document referring to an oral disclosure, use, exhibition or other means																				
"P" document published prior to the international filing date but later than the priority date claimed																				
Date of the actual completion of the international search 12 January 1994		Date of mailing of the international search report 02 FEB 1994																		
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231		Authorized officer KENNETH R. HORLICK <i>K. R. Horlick</i>																		
Facsimile No. NOT APPLICABLE		Telephone No. (703) 305-6196																		

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/10307

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. Immunol., Vol. 141, Number 6, issued 15 September 1988, Gougos et al., "Identification of a human endothelial cell antigen with monoclonal antibody 44G4 produced against a pre-B leukemic cell line", pages 1925-1933, see Fig. 8.	11-12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/10307

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

C07H 21/02, 04; C07K 9/00; A61K 37/00; C12N 1/21, 5/10, 15/70; C12P 21/00

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☒ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**